

STUDY
Mate

ISSUE 8 2008

Mathematics

Mathematical Literacy

English FAL

Afrikaans FAL

Physical Science

Life Sciences

Accounting

Business Studies

History

Geography

GRADE 12

181 DAYS
before final
exams!

Mathematics





National Senior Certificate Grade 12
Mathematics
Exemplar 2008 – Paper 1

MARKS: 136
TIME: 2 hours 43 minutes

- INSTRUCTIONS AND INFORMATION**
Read the following instructions carefully before answering the questions.
- 1. This question paper consists of 11 questions. Answer ALL the questions.
 - 2. Clearly show ALL calculations, diagrams, graphs, *et cetera*, which you have used in determining the answers.
 - 3. An approved scientific calculator (non-programmable and non-graphical) may be used, unless stated otherwise.
 - 4. If necessary, answers should be rounded off to TWO decimal places, unless stated otherwise.
 - 5. Number the answers correctly according to the numbering system used in this question paper.
 - 6. Diagrams are NOT necessarily drawn to scale.
 - 7. It is in your own interest to write legibly and to present the work neatly.
 - 8. A diagram sheet for answering QUESTION 7.3 is included at the end of this question paper. Write your name/examination number in the spaces provided and hand them in together with your ANSWER BOOK.

QUESTION 1

- 1.1 Solve for x :
 - 1.1.1 $x^2 - 10x = 24$ (3)
 - 1.1.2 $x^2 - 6x = 10(1 - 3x)$ (5)
 - 1.1.3 $(x - 1)(x - 2) \leq 6$ (4)
- 1.2 Solve for x and y simultaneously:
 $x + 3y = 5$ and $xy + y^2 = 3$ (7)

QUESTION 2

- 2.1 Determine how long, in years, it will take for the value of a motor vehicle to decrease to 25% of its original value if the rate of depreciation, based on the reducing-balance method, is 21% per annum. (5)

- 2.2 The cost of a bus is R1,2 million. It is expected that the value of this bus will depreciate on a reducing balance per annum to R491 520 in 4 years' time. The price of a new bus is expected to increase by 15% per annum.
 - 2.2.1 Calculate the percentage annual rate of depreciation of the bus. (4)
 - 2.2.2 If the bus needs to be replaced in 4 years' time, calculate the value of the sinking fund that needs to be set up to pay for the new bus. Assume that the old bus will be traded in at its depreciation value of R491 520. (2)
 - 2.2.3 Calculate the percentage increase in the cost of the bus in 4 years. (2)
 - 2.2.4 Calculate the amount that must be invested monthly into a sinking fund to cover the replacement cost of the bus in 4 years' time if the interest paid by the financial institution is 9% per annum compounded monthly. Payments are made at the end of each month. (6)


QUESTION 3

- Consider the following sequence of numbers: 2 ; 5 ; 2 ; 9 ; 2 ; 13 ; 2 ; 17 ; ...
- 3.1 Write down the next TWO terms of the sequence, given that the pattern continues. (2)
 - 3.2 Calculate the sum of the first 100 terms of the sequence. (5)

QUESTION 4


- Consider the sequence: 5 ; 18 ; 37 ; 62 ; 93 ; ...
- 4.1 If the sequence behaves consistently, determine the next TWO terms of the sequence. (2)
 - 4.2 Calculate a formula for the n^{th} term of the sequence. (5)
 - 4.3 Use your formula to calculate n if the n^{th} term in the sequence is 1 278. (4)

QUESTION 5



- In the patterns above each consecutive pattern has more shaded squares than the previous one. The area of the shaded portion of the first pattern is $\frac{1}{4}$ square units. Assume that the pattern behaves consistently, as shown above.
- 5.1 The shaded area in Pattern 2 is $\frac{1}{4} + \frac{3}{16}$. Write down the area of the shaded portions of Patterns 3 and 4. (4)
- 5.2 Write down the area of the shaded portion of the n^{th} pattern in sigma notation. (3)
- 5.3 If the pattern continues without end, what does the area in QUESTION 5.2 approach? (2)

QUESTION 6




- 6.1 Calculate the values of a and b . (2)
- 6.2 Write down the equation of f^{-1} in the form $y = \dots$ (3)
- 6.3 Explain why the inverse of g is not a function. (2)
- 6.4 Write down TWO ways in which the domain of g could be restricted in order that g^{-1} is a function. (2)
- 6.5 Determine the x -values for which: (2)
- 6.5.1 $f^{-1}(x) > 0$ (2)
- 6.5.2 $f(x) - 1 = g(x)$ (2)

QUESTION 7

- Given: $g(x) = \frac{2}{x-3} - 1$
- 7.1 Write down the equations of the asymptotes of g . (2)
- 7.2 Calculate the intercepts of g with the axes. (3)
- 7.3 Use DIAGRAM SHEET 1 to draw the asymptotes and make a neat sketch of g . (3)

QUESTION 8




- 8.1 Write down the period of f . (1)
- 8.2 Give the new range of g if g undergoes a positive vertical shift of 1 unit. (2)
- 8.3 Write down the new equation of f if it is shifted 30° horizontally to the right. (1)

QUESTION 9


- 9.1 Determine $f'(x)$ from first principles if $f(x) = \frac{1}{x}$ (5)
- 9.2 Determine the derivatives of: (2)
- 9.2.1 $f(x) = -5x^2 + 2x$ (2)
- 9.2.2 $y = \sqrt{x^3} + \frac{1}{3x^3}$ (4)

QUESTION 10



- 10.1 Determine the coordinates of A and B. (5)
- 10.2 Determine the turning points of g if $g(x) = f(x - 2)$ (2)
- 10.3 Determine the average rate of change of the function f from A to B. (3)
- 10.4 Determine the equation of the tangent to the graph of f at $x = 1$. (4)
- 10.5 Determine the x -coordinate of the point at which the tangent in QUESTION 10.4 cuts the graph of f again. (4)
- 10.6 Determine the value(s) of k for which $x^3 - 4x^2 - 11x + 30 = k$ will have only one real root. (2)
- 10.7 Determine the point(s) of inflection of f . (6)


QUESTION 11



- The volume of a certain rectangular box is given by the equation $f(x) = x^3 - 8x^2 + 5x + 50$
- 11.1 If the height of the box is $(5 - x)$ units, determine an algebraic expression for the area of the base of the box. (3)
- 11.2 Calculate the value of x for which the volume is a maximum. (6)

DIAGRAM SHEET 1

QUESTION 7





Memorandum

QUESTION 1

1.1.1	$x^2 - 10x = 24$ $x^2 - 10x - 24 = 0$ $(x - 12)(x + 2) = 0$ $x = 12 \quad \text{or} \quad x = -2$ OR $(x - 5)^2 = 49 = 7^2$ $\therefore x - 5 = 7 \quad \text{or} \quad -7$ $\therefore x = 12 \quad \text{or} \quad -2$	✓ standard form ✓ factors ✓ answers	(3)
1.1.2	$x^2 - 6x = 10(1 - 3x)$ $x^2 - 6x = 10 - 30x$ $x^2 + 24x - 10 = 0$ $x = \frac{-24 \pm \sqrt{(24)^2 - 4(1)(-10)}}{2(1)}$ $x = \frac{-24 \pm \sqrt{616}}{2}$ $x = 0,41 \quad \text{or} \quad x = -24,41$	✓ standard form ✓ substitution ✓ 616 ✓✓ answers	(5)
1.1.3	$(x - 1)(x - 2) \leq 6$ $x^2 - 3x - 4 \leq 0$ $(x - 4)(x + 1) \leq 0$ $\begin{array}{ccccccc} + & 0 & & - & & 0 & + \\ \hline & -1 & & & & 4 & \end{array}$ $-1 \leq x \leq 4$	✓ standard form ✓ factors ✓✓ answer	(4)
1.2	$x + 3y = 5$ $x = 5 - 3y$ $(5 - 3y)y + y^2 = 3$ $5y - 3y^2 + y^2 = 3$ $0 = 3 - 5y + 2y^2$ $0 = (3 - 2y)(1 - y)$ $y = \frac{3}{2} \quad \text{or} \quad y = 1$ $x = \frac{1}{2} \quad \text{or} \quad x = 2$	✓ solve for x ✓ substitution ✓ standard form ✓ factors ✓✓ y -answers ✓ x -answers	(7) [19]

QUESTION 2

2.1	$F_v = P_v(1 - i)^n$ $0,25P = P(1 - 0,21)^n$ $n \log 0,79 = \log 0,25$ $n = \frac{\log 0,25}{\log 0,79}$ $n = 5,88 \text{ years}$	✓ formula ✓ substitution ✓ simplification ✓ $n =$ ✓ answer	(5)
2.2.1	$F_v = P_v(1 - i)^n$ $491\,520 = 1\,200\,000(1 - i)^4$ $(1 - i)^4 = 0,409\,6$ $i = 0,2$ $r = 20,00\%$	✓✓ substitution ✓ simplification ✓ answer	(4)
2.2.2	$FV_{(\text{sinkingfund})} = 1,200000(1,15)^4 - 491\,520$ $= R1\,607\,287,50$	✓substitution ✓ answer	(2)
2.2.3	$(1,15)^4 = 1,74900625$ $\therefore \text{an increase of } 74,90\%$	✓ substitution ✓ increase	(2)
2.2.4	Let x be the monthly repayment $i = \frac{9}{1\,200} = 0,007\,5$ $1\,607\,287,50 = x \left[\frac{(1,007\,5)^{48} - 1}{0,007\,5} \right]$ $1\,607\,287,50 = x(57,520\,711\,1)$ $x = R27\,942,76$	✓ i ✓ 1 607 287,50 ✓ 1,0075 ✓ substitution ✓ 57,5207111 ✓ answer	(6)[19]

QUESTION 3

3.1	2 ; 21	✓ 2 ✓ 21	(2)
3.2	$(2 + 2 + \dots + 2) + (5 + 9 + 13 + \dots)$ <i>for 50 terms</i> <i>for 50 terms</i> $= \sum_{i=1}^{50} 2 + \sum_{i=1}^{50} (4i + 1)$ $= 2(50) + \left[\frac{50}{2}(2(5) + 49(4)) \right]$ $= 100 + 25(10 + 196)$ $= 100 + 5\,150$ $= 5\,250$	✓ $2 + 2 + \dots + 2$ <i>for 50 terms</i> ✓ $5 + 9 + 13 \dots$ <i>for 50 terms</i> ✓ 100 ✓ substitution ✓ answer	(5) [7]

QUESTION 4

4.1	130 ; 173	✓ ✓	(2)
4.2	5 18 37 62 93 13 19 25 31 6 6 6 The second difference is constant $\therefore T_n$ is quadratic $\therefore an^2 + bn + c = T_n$$2a = 6$$a = 3$$T_n = 3n^2 + bn + c$$5 = 3(1)^2 + b(1) + c$$b + c = 2 \dots \text{(i)}$$18 = 3(2)^2 + b(2) + c$$2b + c = 6 \dots \text{(ii)}$ (ii) - (i):$b = 4$$c = -2$$T_n = 3n^2 + 4n - 2$ OR $a(1)^2 + b(1) + c = 5$ (i) $a + b + c = 5$ $a(2)^2 + b(2) + c = 18$ (ii) $4a + 2b + c = 18$ $a(3)^2 + b(3) + c = 37$ (iii) $9a + 3b + c = 37$ (ii) - (i):$3a + b = 13$$b = 13 - 3a$ Substitute $b = 13 - 3a$ into (iii)$9a + 3(13 - 3a) + c = 37$$9a + 39 - 9a + c = 37$$c = -2$ Substitute $b = 13 - 3a$ and $c = -2$ into (ii)$4a + 2(13 - 3a) + (-2) = 18$$-2a = -6$$a = 3$$\therefore b = 4$ OR Let T_n be the n^{th} term of the sequence Then $T_2 - T_1 = 13$$T_3 - T_2 = 19$$T_4 - T_3 = 25$$T_5 - T_4 = 31$$T_n - T_{n-1} = \dots$ } add both sides $T_n - T_1 = 13 + 19 + 25 + \dots \text{ (to } n - 1 \text{ terms)}$$T_n - 5 = \left(\frac{n-1}{2}\right)[2(13) + (n-2)6]$$T_n = (n-1)(3n+7) + 5$$T_n = 3n^2 + 4n - 2$	✓ $a = 3$ ✓ substitution ✓ equation (ii) ✓ $b = 4$ ✓ $c = -2$ ✓ subst. $x = 1, 2, 3$ ✓ Equations (ii) - (i): ✓ Substitution into (iii) ✓ $c = -2$ ✓ $b = 4$ ✓ Let T_n ✓ subtracting terms ✓ $T_n - T$ ✓ $T_n - 5$ ✓ $T_n - 3n^2 + 4n - 2$	(5)
4.3	$3n^2 + 4n - 2 = 1\,278$ $3n^2 + 4n - 1\,280 = 0$ $(3n + 64)(n - 20) = 0$ $n = \frac{-64}{3} \quad \text{or} \quad n = 20$ $n = \frac{-64}{3} \text{ is not valid}$ $\therefore n = 20$	✓ substitution ✓ factors ✓ $n \neq \frac{-64}{3}$ ✓ $n = 20$	(4) [11]

QUESTION 5


5.1	Pattern 3 $\frac{1}{4} + \frac{3}{16} + \frac{9}{64}$ Pattern 4 $\frac{1}{4} + \frac{3}{16} + \frac{9}{64} + \frac{27}{256}$	✓ sum ✓ $\frac{9}{64}$ ✓ sum ✓ $\frac{27}{256}$	(4)
5.2	$\frac{1}{4} + \frac{3}{16} + \frac{9}{64} + \dots + \frac{3^{n-1}}{4^n}$ $\sum_{k=1}^n \frac{3^{k-1}}{4^k}$	✓✓ n^{th} term ✓ sigma notation	(3)
5.3	$a = \frac{1}{4} \qquad r = \frac{3}{4}$ $S = \frac{a}{1-r} = \frac{\frac{1}{4}}{1-\frac{3}{4}} = 1$	✓ r ✓ answer	(2) [9]

QUESTION 6

6.1	$b(1)^2 = \frac{1}{2}$ $b = \frac{1}{2}$ $a^1 = \frac{1}{2}$ $a = \frac{1}{2}$	✓ answer ✓ answer	(2)
-----	--	------------------------------	-----

6.2	$y = 2^{-x}$ $y = \left(\frac{1}{2}\right)^x$ $f^{-1}: x = \left(\frac{1}{2}\right)^y$ $y = \log_{\frac{1}{2}} x$	✓ substitution ✓ $f^{-1}: x = \left(\frac{1}{2}\right)^y$ ✓ answer (3)
6.3	The inverse is not a function because for example, $g^{-1}\left(\frac{1}{2}\right) = 1$ or -1	✓✓ answer (2)
6.4	$x \in [0; \infty)$ or $x \in (-\infty; 0]$	✓ $x \in [0; \infty)$ ✓ $x \in (-\infty; 0]$ (2)
6.5.1	$0 < x < 1$	✓✓ answer (2)
6.5.2	$f(x) - 1 = g(x)$ $f(x) - g(x) = 1$ $x = 0$	✓ statement ✓ answer (2) [13]

QUESTION 7

7.1	$x = 3$ $y = -1$	✓ answer ✓ answer (2)
7.2	x -intercept: $0 = -1 + \frac{2}{x-3}$ $x - 3 = 2$ $x = 5$ x -intercept $(5; 0)$ y -intercept $(0; -\frac{5}{3})$	✓ substitution ✓ x -intercept ✓ y -intercept (3)
7.3		✓ shape ✓ intercepts ✓ asymptotes (3) [8]

QUESTION 8


8.1	Period = 360°	✓ answer (1)
8.2	The shift changes the range of g and will now become $[-1; 3]$	✓ -1 ✓ 3 (2)
8.3	$h(x) = \cos(x + 30^\circ - 30^\circ)$ $h(x) = \cos x$	✓ answer (1) [4]

QUESTION 9

9.1	$f'(x) = \lim_{h \rightarrow 0} \frac{\frac{1}{x+h} - \frac{1}{x}}{h}$ $= \lim_{h \rightarrow 0} \frac{\frac{x - (x+h)}{x(x+h)}}{h}$ $= \lim_{h \rightarrow 0} \frac{x - (x+h)}{x(x+h)} \div h$ $= \lim_{h \rightarrow 0} \frac{-h}{x(x+h)} \times \frac{1}{h}$ $= \lim_{h \rightarrow 0} \frac{-1}{x(x+h)}$ $= \frac{-1}{x^2}$	✓ substitution ✓ common denominator ✓ simplification ✓ simplification ✓ answer (5)
9.2.1	$D_x [-5x^2 + 2x]$ $= -10x + 2$	✓✓ answer (2)

9.2.2	$y = \sqrt{x^3} + \frac{1}{3x^3}$ $y = x^{\frac{3}{2}} + \frac{1}{3}x^{-3}$ $\frac{dy}{dx} = \frac{3}{2}x^{\frac{1}{2}} - x^{-4}$	✓ $x^{\frac{3}{2}}$ ✓ $\frac{1}{3}x^{-3}$ ✓✓ answer (4) [11]
-------	---	---

QUESTION 10

10.1	$f'(x) = 3x^2 - 8x - 11$ $0 = (3x - 11)(x + 1)$ $x = \frac{11}{3}$ or $x = -1$ $A(-1; 36)$ and $B(\frac{11}{3}; -14,81)$ Or Some candidates may know Horner's method $f(\frac{11}{3}) = \left[\left(\left(\frac{11}{3} - 4\right) \times \frac{11}{3}\right) - 11\right] \times \frac{11}{3} + 30$ (this makes calculator work much easier) i.e. $f(x) = ((x - 4)x - 11)x + 30$	✓ derivative ✓ derivative = 0 ✓ factors ✓ x -values ✓ points (5)
10.2	$(1; 36)$ $(\frac{17}{3}; -14,8)$	✓ $x = 1$ ✓ y -values remain the same (2)
10.3	Average rate of change $= \frac{36 - (-14,8)}{-1 - \frac{11}{3}}$ $= \frac{50,8}{-4,6}$ $= -10,89$	✓ formula ✓ substitution ✓ answer (3)
10.4	$f'(1) = 3x^2 - 8x - 1$ $= 3(1)^2 - 8(1) - 1$ $= -16$ $f(1) = 1^3 - 4(1)^2 - 11(1) + 30$ $f(1) = 16$ $y - 16 = -16(x - 1)$ $y = -16x + 32$	✓ $m = -16$ ✓ point $(1; 16)$ ✓ substitution ✓ answer (4)
10.5	$-16x + 32 = x^3 - 4x^2 - 11x + 30$ $0 = x^3 - 4x^2 + 6x - 2$ $0 = (x - 1)(x^2 - 3x + 2)$ $0 = (x - 1)(x - 1)(x - 2)$ $x = 1$ or $x = 2$ the tangent cuts the graph again at $x = 2$	✓ substitution ✓ standard form ✓ factors ✓ answer (4)
10.6	$k > 36$ or $k < -14,81$	✓✓ answers (2)
10.7	$f'(x) = 3x^2 - 8x - 1$ $f''(x) = 6x - 8$ $0 = 6x - 8$ $x = \frac{4}{3}$ $f''\left(\frac{4}{3}\right) < 0$ $f''\left(\frac{4}{3}\right) > 0$  Point of inflection $(\frac{4}{3}, \frac{286}{27})$ or $(1,33; 10,59)$	✓ $f''(x) = 6x - 8$ ✓ $f''(x) = 0$ ✓✓ argument ✓ x -value ✓ y -value (6) [26]

QUESTION 11

11.1	area = $\frac{\text{volume}}{\text{height}}$ $= \frac{x^3 - 8x^2 + 5x + 50}{5 - x}$ $= -x^2 + 3x + 10$	✓ division by $5 - x$ ✓✓ answer (3)
11.2	$f(x) = x^3 - 8x^2 + 5x + 50$ $f'(x) = 3x^2 - 16x + 5$ $0 = 3x^2 - 16x + 5$ $0 = (3x - 1)(x - 5)$ $x = \frac{1}{3}$ or $x \neq 5$ But $x = 5$ is not valid $\therefore x = \frac{1}{3}$	✓ $f'(x)$ ✓ $= 0$ ✓ factors ✓ answers ✓ reject $x = 5$ ✓ dimensions (6) [9]

TOTAL 136